

# 11th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions



Beitrag ID: 296

Typ: Talk

## Data-driven $\hat{q}$ in a hard-soft factorized parton energy loss approach

Mittwoch, 29. März 2023 09:40 (20 Minuten)

Interactions of high-energy partons with the strongly-coupled quark-gluon plasma lead to parton energy loss, as well as broadening of the partons' transverse and longitudinal momentum distributions. Energy loss and momentum broadening resulting from soft parton-plasma interactions can be quantified with transport coefficients, factorizing their effect from hard (perturbative) parton-plasma scatterings. We apply this factorized model of energy loss [1] to perform a Bayesian calibration against RHIC and LHC measurements, finding an enhancement of parton energy loss at low temperature compared to perturbative expectations. We highlight the model's ability to match perturbative calculations while inherently allowing for agnostic non-perturbative energy loss.

[1] T. Dai, J.-F. Paquet, D. Teaney and S.A. Bass, "Parton energy loss in a hard-soft factorized approach," Phys.Rev.C 105 (2022) no.3, 034905

### Experiment/Theory

Theory/Phenomenology

### Affiliation

Duke University;  
Vanderbilt University

**Primary authors:** DAI, Tianyu (Duke University); PAQUET, Jean-Francois (Vanderbilt University); BASS, Steffen A. (Duke University)

**Vortragende(r):** BASS, Steffen A. (Duke University)

**Sitzung Einordnung:** Parallel: Jets and their modification in QCD Matter

**Track Klassifizierung:** Jets and their modification in QCD matter