

Photo-nuclear Jet Production in Ultra-Peripheral Pb+Pb Collisions at 5.02 TeV with the ATLAS Detector


Ben Gilbert

[ATLAS-CONF-2022-021](#)




Introduction: Photo-nuclear Jet Production

In ultra-relativistic heavy ion collisions, the intense electromagnetic fields provide a flux of quasi-real photons.



Introduction: Photo-nuclear Jet Production

In ultra-relativistic heavy ion collisions, the intense electromagnetic fields provide a flux of quasi-real photons.




Those photons can scatter off of nuclear partons.

Introduction: Photo-nuclear Jet Production

In ultra-relativistic heavy ion collisions, the intense electromagnetic fields provide a flux of quasi-real photons.

“Direct” photons scatter off the nucleus.

“Resolved” photons scatter through a virtual hadronic excitation of the photon.





Those photons can scatter off of nuclear partons.

Introduction: Photo-nuclear Jet Production

In ultra-relativistic heavy ion collisions, the intense electromagnetic fields provide a flux of quasi-real photons.

“Direct” photons scatter off the nucleus.

“Resolved” photons scatter through a virtual hadronic excitation of the photon.




Jet kinematics provide access to hard-scattering kinematics, directly probing nuclear PDF effects.

$$H_T = \sum_i p_T^i \quad x_A = \frac{M_{jets} e^{-y_{jets}}}{\sqrt{s_{NN}}} \quad z_\gamma = \frac{M_{jets} e^{+y_{jets}}}{\sqrt{s_{NN}}}$$

Those photons can scatter off of nuclear partons.

Introduction: Nuclear PDFs at Low- x

- Nuclear Parton Distribution Functions (nPDFs) are important for precision measurements of a number of physical observables.
- They are poorly constrained at low- x and intermediate Q^2 due to a lack of available data.
 - $100 \text{ GeV}^2 < Q^2 < 1000 \text{ GeV}^2$ has very little constraint.
 - Nuclear shadowing at low- x in this region is of particular theoretical interest.
- Photo-nuclear jet production provides a clean probe of this kinematic region, similar to DIS:
 - Proposal by [Strikman, Vogt, and White \(2005\)](#)
 - Test of sensitivity (right) by [Helenius \(2018\)](#)



Selecting Photo-nuclear Jet Events

Event Selections


Raw Yields

Unfolding

Systematic
Uncertainty

Final Results

Event Selections



Selecting Photo-nuclear Jet Events

Event Selections

Raw Yields



Unfolding

Systematic
Uncertainty


Final Results

Event Selections

- OnXn requirement for nuclear breakup in exactly one ATLAS Zero-Degree Calorimeter (ZDC)






Selecting Photo-nuclear Jet Events




Event Selections

- OnXn requirement for nuclear breakup in exactly one ATLAS Zero-Degree Calorimeter (ZDC)
- Large rapidity gaps on one side of the detector
 - To veto $\gamma\gamma \rightarrow q\bar{q}$, also require $\Delta\eta_A^{edge} < 3$.






Selecting Photo-nuclear Jet Events




Event Selections

- OnXn requirement for nuclear breakup in exactly one ATLAS Zero-Degree Calorimeter (ZDC)
- Large rapidity gaps on one side of the detector
 - To veto $\gamma\gamma \rightarrow q\bar{q}$, also require $\Delta\eta_A^{edge} < 3$.




Selecting Photo-nuclear Jet Events



Event Selections

- OnXn requirement for nuclear breakup in exactly one ATLAS Zero-Degree Calorimeter (ZDC)
- Large rapidity gaps on one side of the detector
 - To veto $\gamma\gamma \rightarrow q\bar{q}$, also require $\Delta\eta_A^{edge} < 3$.
- At least two Particle-Flow jets with $p_T > 15$ GeV.



Selecting at least two jets allows access to the hard-scattering kinematics.



$$H_T = \sum_i p_T^i$$

$$x_A = \frac{M_{jets} e^{-y_{jets}}}{\sqrt{S_{NN}}}$$

$$z_\gamma = \frac{M_{jets} e^{+y_{jets}}}{\sqrt{S_{NN}}}$$



Importance of Forward Neutrons: $XnXn$ Events

Event Selections


Raw Yields

Unfolding

Systematic
Uncertainty

Final Results

The photo-nuclear jet requirements select events with very high-energy photons.



Importance of Forward Neutrons: $XnXn$ Events

Event Selections

Raw Yields



Unfolding

Systematic
Uncertainty

Final Results

The photo-nuclear jet requirements select events with very high-energy photons.

- $E_\gamma \propto 1/b \rightarrow$ Biases towards lower impact parameter collisions
- Much higher probability of breakup due to additional EM interactions



Importance of Forward Neutrons: $XnXn$ Events

Event Selections

Raw Yields


Unfolding

Systematic
Uncertainty


Final Results

The photo-nuclear jet requirements select events with very high-energy photons.


- $E_\gamma \propto 1/b \rightarrow$ Biases towards lower impact parameter collisions
- Much higher probability of breakup due to additional EM interactions



Studies of dijet events with large gaps on one side estimate about 50% of photo-nuclear jet production breaks up!




Importance of Forward Neutrons: $XnXn$ Events



This theoretical model for breakup is used to compare theory to data.


The photo-nuclear jet requirements select events with very high-energy photons.

- $E_\gamma \propto 1/b \rightarrow$ Biases towards lower impact parameter collisions
- Much higher probability of breakup due to additional EM interactions



Studies of dijet events with large gaps on one side estimate about 50% of photo-nuclear jet production breaks up!

Basic theoretical modeling predicts an even higher rate.




Importance of Forward Neutrons: $0n0n$ Events

ATLAS has observed inclusive jet production in UPCs without nuclear breakup ($0n0n$).

Gaps are required on both sides of the detector: $\sum \Delta\eta > 2.0$

A factor of 10 more events are observed in data than are predicted from $\gamma\gamma \rightarrow$ jets, estimated by Pythia or comparison to $\gamma\gamma \rightarrow \mu^+\mu^-$ studies.





Importance of Forward Neutrons: $0n0n$ Events

ATLAS has observed inclusive jet production in UPCs without nuclear breakup ($0n0n$).

Gaps are required on both sides of the detector: $\sum \Delta\eta > 2.0$

A factor of 10 more events are observed in data than are predicted from $\gamma\gamma \rightarrow$ jets, estimated by Pythia or comparison to $\gamma\gamma \rightarrow \mu^+\mu^-$ studies.

The distribution shapes are clearly different from pure $\gamma\gamma \rightarrow$ jets.




Importance of Forward Neutrons: $0n0n$ Events

ATLAS has observed inclusive jet production in UPCs without nuclear breakup ($0n0n$).

Gaps are required on both sides of the detector: $\sum \Delta\eta > 2.0$



A factor of 10 more events are observed in data than are predicted from $\gamma\gamma \rightarrow$ jets, estimated by Pythia or comparison to $\gamma\gamma \rightarrow \mu^+\mu^-$ studies.

The distribution shapes are clearly different from pure $\gamma\gamma \rightarrow$ jets.



Constructing the Cross-Section


$$\frac{d^3\sigma}{dH_T dx_A dz_\gamma} = \frac{1}{\mathcal{L}} \frac{\Delta Y}{\Delta H_T \Delta x_A \Delta z_\gamma}$$



$$H_T = \sum_i p_T^i$$


$$x_A = \frac{M_{jets} e^{-y_{jets}}}{\sqrt{s_{NN}}}$$

$$z_\gamma = \frac{M_{jets} e^{+y_{jets}}}{\sqrt{s_{NN}}}$$




Constructing the Cross-Section

$$\frac{d^3\sigma}{dH_T dx_A dz_\gamma} = \frac{1}{\mathcal{L}} \frac{\Delta Y}{\Delta H_T \Delta x_A \Delta z_\gamma}$$



H_T does not depend strongly on x_A or z_γ .




$$H_T = \sum_i p_T^i$$


$$x_A = \frac{M_{jets} e^{-y_{jets}}}{\sqrt{s_{NN}}}$$

$$z_\gamma = \frac{M_{jets} e^{+y_{jets}}}{\sqrt{s_{NN}}}$$

Acceptance in x_A is strongly dependent on the photon energy, z_γ .



Unfolding Measured Cross-Sections





$$H_T = \sum_i p_T^i$$

$$x_A = \frac{M_{jets} e^{-y_{jets}}}{\sqrt{s_{NN}}}$$

$$z_\gamma = \frac{M_{jets} e^{+y_{jets}}}{\sqrt{s_{NN}}}$$

The measured cross-sections are then unfolded in 3 dimensions to correct for detector effects.

- Low- p_T flavor effects are the largest correction.



Systematic Uncertainties

Event Selections

Raw Yields

Unfolding

Systematic
Uncertainty


Final Results

Systematic uncertainties are the key limiting factor in our sensitivity to nuclear PDFs.

The jet energy **scale** and **resolution** uncertainties are typically 5-10%.

Control over the **preliminary low- μ calibration** currently provides the dominant source of uncertainty.

Systematic uncertainties are also evaluated on the **unfolding** and **event selections**.



Systematic Uncertainties

Event Selections

Raw Yields

Unfolding

Systematic
Uncertainty


Final Results

Systematic uncertainties are the key limiting factor in our sensitivity to nuclear PDFs.

The jet energy **scale** and **resolution** uncertainties are typically 5-10%.

Control over the **preliminary low- μ calibration** currently provides the dominant source of uncertainty.

Systematic uncertainties are also evaluated on the **unfolding** and **event selections**.



The Measured Photon Flux

Event Selections



Raw Yields

Unfolding

Systematic
Uncertainty

Final Results

- The distribution of z_γ values for large x_A in bins of H_T (right) demonstrates the measured photon flux.
 - The breakup model performs well within systematic uncertainties.
 - Disagreements appear to arise at low z_γ , where the breakup model tends to over-correct.



Scanning in Photon Energy

Event Selections

Raw Yields

Unfolding


Systematic
Uncertainty

Final Results

$$H_T = \sum_i p_T^i$$


$$x_A = \frac{M_{jets} e^{-y_{jets}}}{\sqrt{S_{NN}}}$$

$$z_\gamma = \frac{M_{jets} e^{+y_{jets}}}{\sqrt{S_{NN}}}$$



The x_A distribution has substantial acceptance effects in z_γ .

Selecting on photon energy removes this bias, allowing a more direct measurement of nPDFs.



Scanning in Photon Energy

Event Selections

Raw Yields

Unfolding


Systematic
Uncertainty

Final Results

$$H_T = \sum_i p_T^i$$


$$x_A = \frac{M_{jets} e^{-y_{jets}}}{\sqrt{S_{NN}}}$$

$$z_\gamma = \frac{M_{jets} e^{+y_{jets}}}{\sqrt{S_{NN}}}$$



The x_A distribution has substantial acceptance effects in z_γ .

Selecting on photon energy removes this bias, allowing a more direct measurement of nPDFs.



Measured Cross-Sections

Event Selections


Raw Yields

Unfolding


Systematic
Uncertainty

Final Results

- At intermediate photon energies, we can access higher-x partons.



Photon Energy
 0.004 < z_γ < 0.008



Measured Cross-Sections

Event Selections


Raw Yields

Unfolding


Systematic
Uncertainty

Final Results


- Higher photon energy opens up the low-x shadowing region.
- Results are quite consistent with the theoretical model.




Photon Energy
0.008 < z_γ < 0.015




Measured Cross-Sections



- The highest photon energy allows the most access to low x .
- Systematic control is more challenging near acceptance edges.




Photon Energy
 0.015 < z_γ < 0.027



Conclusions and Next Steps

- Photo-nuclear jet production was measured by ATLAS in 5.02 TeV Pb+Pb collisions with 2018 data.
 - Particle-Flow jets allow for the measurement to be extended even lower in jet p_T while maintaining systematic control.
 - This measurement has been fully unfolded for detector response for the first time.
- The overall normalization of the cross-section is well-predicted by theoretical comparisons.
 - A theoretical model of nuclear breakup is necessary to understand the total cross-section.
- This study is sensitive to nuclear PDF effects with a precision of up to 10% in some bins.
 - Once final studies of low- μ jet response in ATLAS can be completed, substantial gains in systematic control are possible.
- These results are connected to early physics goals for the EIC.



Backup

Theoretical Modeling of Nuclear Breakup

- The photon flux available through Pythia makes certain overly-simplified assumptions which we correct via modeling with STARlight.

We integrate over A-A impact parameter (b) and the impact parameter relative to the photon-emitting nucleus (s_A).

Correction for the probability of breakup due to additional EM interactions

Nuclear thickness function

$$F_{\gamma/A}^{\text{eff}}(E_\gamma) \equiv \int \underbrace{d^2b}_{\text{black}} \underbrace{d^2s_A}_{\text{red}} \underbrace{P_{\text{no had}}(b)}_{\text{red}} \underbrace{P_{\text{no EM}}(b)}_{\text{blue}} \underbrace{f_{\gamma/A}(E_\gamma, s)}_{\text{magenta}} \underbrace{T_B(\vec{s}_A - \vec{b})}_{\text{green}}$$

Correction for the probability of breakup due to hadronic interactions (overlap veto)

The photon flux from Pythia uses a point source, so this term corrects for coherent nuclear emission.

Measured Cross-Sections

Event Selections


Raw Yields

Unfolding


Systematic
Uncertainty

Final Results

- At intermediate photon energies, we can access higher-x partons.



Photon Energy
0.004 < z_γ < 0.008
👉



Measured Cross-Sections

Event Selections


Raw Yields

Unfolding


Systematic
Uncertainty

Final Results

- Higher photon energy opens up the low-x shadowing region.
- Results are quite consistent with the theoretical model.



Photon Energy
0.008 < z_γ < 0.015



Measured Cross-Sections

Event Selections


Raw Yields

Unfolding

Systematic
Uncertainty

Final Results

- The highest photon energy allows the most access to low x .
- Systematic control is more challenging near acceptance edges.



Photon Energy
 $0.015 < z_\gamma < 0.027$

