

11th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions



Beitrag ID: 120

Typ: Poster

Longitudinal momentum fraction of heavy flavor meson in jets in high-energy nuclear collisions

Dienstag, 28. März 2023 18:15 (2 Stunden)

Heavy flavor jets are powerful tools to gain insight into the in-medium partonic energy loss mechanisms and the transport properties of the quark-gluon plasma (QGP) in high-energy nuclear collisions. In this work, we present the first theoretical study of the longitudinal momentum fraction $z_{||}$ carried by D^0 meson in jets in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. The p+p baseline is provided by POWHEG+PYTHIA8 which matches the next-to-leading order hard processes with the parton shower. The in-medium evolution of heavy quark jets is employed by a Monte Carlo transport model which takes into account the collisional and radiative partonic energy loss in the expanding QGP. We observe steeper $z_{||}$ distributions of B^0 -jet compared to that of D^0 -jet at the same kinematics region in p+p collisions, which may be a hint of the harder jet fragmentation function of b-jet compared to c-jet in vacuum. In A+A collisions, it is shown that the jet quenching effect would in general decrease the values of $z_{||}$. In addition, we predict visibly stronger nuclear modifications of B^0 -jet $z_{||}$ distributions compared to D^0 -jet within the same p_T windows, as a result of the much steeper initial $z_{||}$ distribution of B^0 -jet in vacuum.

Experiment/Theory

Theory/Phenomenology

Affiliation

Central China Normal University

Primary authors: LI, Yao (Central China Normal University); DAI, Wei (China university of Geosciences); ZHANG, Ben-Wei (Central China Normal University); Dr. WANG, Sa (South China Normal University)

Vortragende(r): DAI, Wei (China university of Geosciences)

Sitzung Einordnung: Poster Session

Track Klassifizierung: Heavy flavor and quarkonia