


# Future Facilities: The EIC

**The ePIC detector -  
From physics motivation  
to a viable detector concept**

**Friederike Bock (ORNL)  
March 30, 2023**



**HP 2023**, Aschaffenburg, Germany




# Back to the basics!

## Where we are:


- Elastic lepton scattering determined the nucleon's charge & magnetism distributions in sphere with  $\langle r_{ch} \rangle \approx 0.84$  fm
- Large fraction of momentum in proton ( $x$ ) carried by 3 valence quarks (2u,d), but very small fraction of proton spin
- Nucleons contain additional dynamically generated quark-antiquark pairs & gluons each carrying low fraction of momentum
- Quark & gluon longitudinal momentum fractions well mapped out
- Nucleon spin & mass have large contributions from quark-gluon dynamics, described by QCD




Proton  
early 1900s





Proton  
1975



Proton  
2015




Proton  
in a nucleus



# How did we learn this?



## Deep Inelastic Scattering (DIS)



$$Q^2 = s \cdot x \cdot y$$


$s$  center-of-mass energy squared

$Q^2$  resolution power

$x$  the fraction of the nucleon's momentum carried by the struck quarks ( $0 < x < 1$ )


$y$  inelasticity

- As a probe, electron beams provide unmatched precision of the electromagnetic interaction
- Event-by-event, model independent leading order determination of parton kinematics including sensitivity to particle's spin is possible
- Data at higher  $Q^2$  obtained indirectly from hadron-collider measurements



# What we don't know yet


- The 3D distributions of sea quarks & gluons and their spins in nucleon
- How do the **nucleon mass & spin** emerge from them and their interactions?
- The details of **interactions of color-charged quarks and gluons with a nuclear medium**
- How are **nuclear bindings and hadronic states** created from quark, gluons and their interactions?
- How does a dense nuclear environment affect the quarks and gluons and their interactions?
- The **gluon density in nuclei**
- Is there a **Color Glass Condensate**?



# EIC vs HERA

## Machine parameters

- **Center-of-mass energy:** 20 - 140 (318) GeV
  - ▶ electrons: 2.5 - 18 (27.5) GeV
  - ▶ protons: 40- 275 (920) GeV (ions:  $Z/A \times E_p$ )
- **Luminosity:**  $10^{34}$  ( $10^{31}$ )  $\text{cm}^{-2} \text{s}^{-1}$
- **Polarization:** up to 70% (e & ion) (only  $e^\pm$  up to 60%)
- **Ion species:**  $p \rightarrow U$  ( $A > 1$  only in fixed target)
- **Detectors:**
  - ▶ full coverage: 2 (2)
  - ▶ fixed target: 0 (2 - limited far-forward coverage)




[EIC CDR]

**Start date:** ~ 2031  
**Location:** BNL  
**Budget:** ~ \$2.4 billion

## EIC will have:


- lower energy
- variable C-o-M energy w/o significant loss in luminosity
- higher luminosity

- + Hadron polarization
- + Nuclear beams
- + Modern detector(s)




[HERA proposal]

# Generalized detector design considerations



- Large rapidity coverage for central detector
- Specialized far-forward detectors for  $p$  kinematics measurements
- High precision low mass tracking
- Hermetic coverage of tracking, electromagnetic & hadronic calorimetry
- High performance single track PID for  $\pi$ ,  $K$ ,  $p$  separation


- Large acceptance for diffraction, tagging, neutrons from nuclear breakup  
many auxillary detectors integrated in beam line: low- $Q^2$  tagger, Roman pots, ZDCs ...
- High control of systematics  
luminosity monitors, electron & hadron polarimetry




## Highly integrated design between detector and machine for IR

# The detector design process


## Define physics objectives & generic design parameters




2012



2015




2017




2020

## Realistic machine & detector concepts



Feb. 2021




Dec. 2021




- Detector & machine design driven by physics objectives
- Jan. 2020: BNL site selection
- Extensive generic detector R&D for EIC for PID, tracking & calorimetry
- YR outlines general detector requirements for benchmark physics observables
- Mar. 2021: Call for Detector Proposals
- Mar. 2022: ECCE chosen as reference design for detector 1
- Jul. 2022: ePIC collaboration
- now: TDR preparations

# The detector design process


## Define physics objectives & generic design parameters




2012



2015




2017




2020

## Realistic machine & detector concepts




Feb. 2021



Dec. 2021




- Detector & machine design driven by physics objectives
- Jan. 2020: BNL site selection
- Extensive generic detector R&D for EIC for PID, tracking & calorimetry
- YR outlines general detector requirements for benchmark physics observables
- Mar. 2021: Call for Detector Proposals
- Mar. 2022: ECCE chosen as reference design for detector 1
- Jul. 2022: ePIC collaboration
- now: TDR preparations





# Tracker layout



## Technology mix

- **ITS3 MAPS based Si-detectors:**  
 $\sigma = 10\mu\text{m}$ ,  $X/X_0 \sim 0.05 - 0.55\%/layer$
- **Gaseous tracker:**  
 $\sigma = 55\mu\text{m}$ ,  $X/X_0 \sim 0.2\%/layer$
- **AC-LGADs:**  
 $\sigma = 30\mu\text{m}$ ,  $X/X_0 \sim 1.5 - 6\%/layer$

- **mid-rapidity:**



Ultra thin MAPS based Si-detectors, gaseous detectors & AC-LGADs

- **Forward and Backward:**


MAPS based Silicon discs & AC-LGADs

- Outer layers placed to provide seeds for tracking & ideal track points before/after PID detectors


- **New Magnet with BABAR dimensions**  $B = 1.7\text{-}2\text{T}$



# Tracking performance



- Stringent requirements from Yellow Report for electron resolution
- Backward momentum resolution requirement hard to meet, complemented by calorimetric resolution
- YR requirement assumes Calorimetry & Tracking need to fulfill requirements independently
- Rapidly evolving tracker design, including background and pattern recognition



E. Yeats, R. Cruz-Torres, N. Schmidt, S. Maple




# Cherenkov-PID

**pfRICH**

**mRICH**

**hpDIRC**

**dRICH**




- Optimized for charged pion, kaon & proton separation
- Particular focus on large  $\eta$  coverage
- Complemented by calorimetry & TOF
- Geometries optimized to fit ECCE baseline design while maintaining required performance
- Two alternatives for backward region, pfRICH & mRICH


⇒ Global optimization process ongoing

# Cherenkov-PID


**pfRICH**




**mRICH**







**hpDIRC**




**dRICH**





**Review 20.-21.3.  
decision anticipated soon**



# Time of flight (TOF)





- Analog Coupled - Low Gain Avalanche Detectors (AC-LGADs) with 25 ps time resolution resolution
- Combined PID & tracking detector
- Positions optimized for low momentum  $e/\pi$ ,  $\pi/K$ ,  $K/p$  separation
- Full  $\eta$ -coverage for simultaneous start time determination
- Alternative barrel design with less  $X/X_0$



O. Hartbrich, N. Schmidt, Z. Ye




# Electromagnetic Calorimetry (1)



## Optimization criteria

- Minimal acceptance gaps
- Scattered electron detection & identification (energy resolution &  $E/p$ )
- Shower separation within jets & good energy resolution (h-endcap)
- Most stringent constraints in e-endcap & barrel
- h-endcap with high granularity & good energy resolution

# Electromagnetic Calorimetry (2)



**Review 13.-14.3.**

**decision anticipated soon**




## Endcap regions:

- **EEMC** - homogenous high resolution  $\text{PbWO}_4$  crystal ECal
- **FEMC** - highly granular W-Scintillating Fiber calorimeter

## Barrel region - alternatives:

- **Sci-Glass**: homogenous, projective Sci-Glass ECal
- **Imaging**: 6 layers of  $0.5 \times 0.5$  mm Astro-Pix Silicon layers, interleaved with Pb-SciFi calorimeter

- Designed to complement tracking in Particle-Flow algorithm
- OHCAL/IHCAL**
  - Fe/Scint sampling calorimeter
  - partial sPHENIX re-use & magnet flux return
- LFHCAL**
  - Fe/Scint & W/Scint sampling calorimeter
  - Highly segmented (7 long. segments)
  - W-segment as colimator
- High granularity inserts under discussion for forward E&HCAL to extend  $\eta$  coverage to  $\eta = 4$
- Electron end-cap HCal as neutral veto, shallow Fe/Scint calo




|              | Barrel HCal                  | LFHCAL                       |
|--------------|------------------------------|------------------------------|
| $\eta$       | [-1 .. 1]                    | [1 .. 4]                     |
| $\sigma_E/E$ | $\sim 75\%/\sqrt{E} + 15\%*$ | $\sim 43\% \sqrt{E} + 5.5\%$ |
| depth        | $\sim 4-5 \lambda_1$         | $\sim 7-8 \lambda_1$         |

\*Based on prototype beam tests and earlier experiments



# Far-forward Region



- **B0 system** for charged-particle measurement in forward direction & neutral-particle tagging
- **off-momentum detectors** measure charged particles with different rigidity than the beam, e.g., those following decay and fission.
- **roman pot detectors** charged particles measurement close to beam envelope
- **zero-degree calorimeter** measures neutral particles at small angles.

| Detector                            | Acceptance   |
|-------------------------------------|--|
| Zero-Degree Calorimeter (ZDC)       | $\theta < 5.5 \text{ mrad } (\eta > 6)$                |
| Roman Pots (2 stations)             | $0 < \theta < 5.0 \text{ mrad } (\eta > 6)$            |
| Off-Momentum Detectors (2 stations) | $\theta < 5.0 \text{ mrad } (\eta > 6)$                |
| B0 detector                         | $5.5 < \theta < 20.0 \text{ mrad } (4.6 < \eta < 5.9)$ |




# Far-backward Region

## Luminosity Monitor

$$e + p \rightarrow e \gamma p$$

$$e + Au \rightarrow e \gamma Au$$

AC-LGAD and PbWO<sub>4</sub> ECal to provide  $\delta L/L \sim 1\%$  or rel.  $L$  determination exceeding  $10^{-4}$  precision



## Low Q<sup>2</sup>-tagger

clean photo-production signal for  $10^{-3} < Q^2 < 10^{-1}$


- Double-layer AC-LGAD tracker at 24 & 37m from IP
- PbWO<sub>4</sub> ECal (20cm x 2cm<sup>2</sup> crystals)

- This area is designed to measure scattered electrons at small, far-backward angles
- Strong technology synergies with central detector systems

# How to access partons at EIC



New Simulations

## Neutral current (SI)DIS




## Neutral current (SI)DIS

- Detect scattered lepton (DIS) in coincidence with identified hadrons (SIDIS)
  - ▶ measure correlation between different hadrons as fct. of  $p_T, z, \eta$
  - ▶ needs FF to correlate hadron type with parton




Ralf Seidl

## Jets



## Charged current DIS



## Charged current DIS - W-exchange

- direct access to the quark flavor no FF – complementary to SIDIS



## Jets

- best observable to access parton kinematics
- tag partons through the sub-processes and jet substructure
  - ▶ di-jets: relative  $p_T \rightarrow$  correlated to  $k_T$
  - ▶ tag on PDF

# Nucleon Spin

$$\frac{1}{2}\hbar = \left\langle P, \frac{1}{2} \left| J_{QCD}^z \right| P, \frac{1}{2} \right\rangle = \underbrace{\frac{1}{2} \int_0^1 dx \Delta\Sigma(x, Q^2)}_{\text{total quark spin}} + \underbrace{\int_0^1 dx \Delta G(x, Q^2)}_{\text{gluon spin}} + \underbrace{\int_0^1 dx \left( \sum_q L_q^z + L_g^z \right)}_{\text{angular momentum}}$$

- quark contribution: integral of  $g_1$  over  $x$  from 0 to 1
- gluon contribution:  $dg_1(x, Q^2)/d\ln Q^2 \rightarrow \Delta g(x, Q^2)$
- Measured through DIS cross section asymmetry in oppositely polarized collisions
- Improved constraints on the spin of quarks/gluons  
 $\Rightarrow$  Constrain contribution of orbital angular momentum (OAM) of partons to proton spin
- Collisions with polarized deuterons/helium-3  
 $\Rightarrow$  Access to neutron spin




New Simulations

Charlotte v. Hulse

# Imaging Nuclei

## Diffraction vector meson production



### DGLAP




- predicts  $Q^2$  but not  $A$ -dependence and  $x$ -dependence

### Saturation models

- predict  $A$ -dependence and  $x$ -dependence but not  $Q^2$

### Need: large $Q^2$ lever-arm for fixed $x$ , $A$ -scan


- Measure different structure function in  $eA \rightarrow$  constrain nPDF
- Does the nucleus behave like a proton at low- $x$ ?
- Diffractive  $J/\psi$  production for imaging nucleus
- Diffractive  $\phi/\rho$  production as saturation probe



New Simulations

# Imaging Nuclei

## Diffraction vector meson production



### DGLAP




- predicts  $Q^2$  but not  $A$ -dependence and  $x$ -dependence

### Saturation models


- predict  $A$ -dependence and  $x$ -dependence but not  $Q^2$

### Need: large $Q^2$ lever-arm for fixed $x$ , $A$ -scan

- Measure different structure function in  $eA \rightarrow$  constrain nPDF
- Does the nucleus behave like a proton at low- $x$ ?
- Diffractive  $J/\psi$  production for imaging nucleus
- Diffractive  $\phi/\rho$  production as saturation probe




New Simulations




# The EIC is coming fast!

## Exiting times ahead! Join us!





# Thank you!

# Back-up





# 2+1 dimensional Imaging of Quarks & Gluons




## Nuclear Femtography

- Structure mapped in terms of:
  - $b_T$  = transverse position
  - $k_T$  = transverse momentum
- use different processes to access different aspects of distribution functions

- **PDFs:** (SI)DIS cross sections
- **GPDs:** Deep Exclusive Scattering (DES) cross sections like: deeply virtual Compton scattering (DVCS)  $\gamma^* + p \rightarrow \gamma + p$  or production of a vector meson  $\gamma^* + p \rightarrow V + p$   
Spin-dependent 2+1D coordinate space images
- **TMDs:** SIDIS cross sections  
Spin-dependent 3D momentum space images


# Color Glass Condensate?



- $e$  interacts over distances  $L \sim (2mN\lambda)^{-1}$
  - For  $L > 2R_A \sim A^{1/3}$  probe cannot distinguish between nucleons in front or back
  - Probe interacts coherently with all nucleons
- ⇒ **Enhancement of  $Q_s$  with  $A \rightarrow$  non-linear QCD regime reached at significantly lower energy in  $A$  than in proton**



## Di-Hadron or Di-Jet Correlations

- Low  $p/A$  gluon density (ep): pQCD  $2 \rightarrow 2$  process predicts  $\Rightarrow$  back-to-back di-jet
- High gluon density (eA):  $2 \rightarrow$  many process  $\Rightarrow$  expect broadening of away-side
- **EIC allows to study the evolution of  $Q_s$  with  $x$**




# Kinematic Coverage

## Collins & Sivers




## DVCS




## polarized DIS

[EIC YR]



**Accelerator gives access to extensive kinematic range**

**⇒ Now we need a detector to match**



**design luminosity:**

$$L = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$$

$$\int L dt = 100 \text{ fb}^{-1} \text{ per year}$$

$$\int L dt \quad \mathbf{1 \text{ fb}^{-1}}$$

**inclusive DIS**

- measure scattered electron
- precision EM-Calorimetry
- multi-dimensional binning:  $x, Q^2$
- maximize  $x, Q^2$  coverage & determines interaction region design

$$\mathbf{10 \text{ fb}^{-1}}$$

**semi-inclusive DIS**

- measure scattered electron in coincidence with identified hadrons
- multi-dimensional binning:  $x, Q^2, z, \theta, p_T$
- maximize PID detector coverage in whole phase space

$$\mathbf{10-100 \text{ fb}^{-1}}$$

**Exclusive processes**

- measure all particles in event
- hermetic tracking + hadronic calorimetry
- multi-dimensional binning:  $x, Q^2, z, \theta, p_T$
- measure proton kinematics
- strong constraints on far-forward detector & interaction region

# Diffractive Vector Meson Production

