


The Quark-Gluon Plasma

Characterizing the early Universe matter with the first 10 years of heavy-ion collisions at the LHC

Anton Andronic




Outline

- Introduction, theoretical basis
- The LHC and the detectors; what is measured
- Hadrons with light-flavor (u,d,s) and the QCD phase diagram
- Collective flow, hydrodynamics
- Jet quenching (parton energy loss)
- Heavy quarks and deconfined matter
- Summary

see reviews: AA, [arXiv:1407.5003](https://arxiv.org/abs/1407.5003); Busza, Rajagopal, van der Schee, [1802.04801](https://arxiv.org/abs/1802.04801)

The proton at high resolution (energies)

A. Andr



3

There are more quarks than the “constituent” (valence) u, u, d
...virtual quark-antiquark pairs; also
many gluons

This is one of many possible configurations
...obtained in (deep-inelastic) electron scattering experiments with resolution 10^{-18} - 10^{-19} m (leading to “parton distribution functions”)

The quarks are never found free
quark confinement (principle)

...are free only asymptotically ($Q \rightarrow \infty$)
Gross, Wilczek, Politzer, 1974 (Nobel Prize 2004)

...has led to the proposal of the Quark-Gluon Plasma (high T)


Collins & Perry, Cabibbo & Parisi, 1975 (Itoh, 1970; Carruthers, 1973; Shuryak)

Lattice QCD predicts a phase transition ($\mu_B=0$)

A. Andronic

numerical solutions of QCD on a discrete (space-time) grid (sophisticated formalism, huge computers)

4




Borsanyi et al., PLB 370 (2014) 99 , PRL 125 (2020) 052001 ; Bazavov et al., PLB 795 (2019) 15

$T_c=156.5 \pm 1.5$ (158.0 \pm 0.6) MeV, $\varepsilon_c \simeq 0.4$ GeV/fm³, or $2.5\varepsilon_{nuclear}$

ideal-gas limit not reached at very large temperatures (GeV range explored)


HISTORY OF THE UNIVERSE



How to "simulate" in laboratory the early Universe?

A. Andronic

6




$$t \simeq 10^{-23} \text{ s}, V \simeq 10^{-40} \text{ m}^3$$

1. initial collisions ($t \leq t_{coll} = 2R/\gamma_{\text{cm}}c$; $R_{Pb} \simeq 7 \text{ fm}$)
2. thermalization: equilibrium is established ($t \lesssim 1 \text{ fm}/c = 3 \times 10^{-24} \text{ s}$)
3. expansion ($\sim 0.6c$) and cooling ($t < 10-15 \text{ fm}/c$) ...deconfined stage?
4. hadronization (quarks and gluons form hadrons)
5. chemical freeze-out: inelastic collisions cease; particle identities (yields) frozen
6. kinetic freeze-out: elastic collisions cease; spectra are frozen ($t+ = 3-5 \text{ fm}/c$)

The CERN accelerator complex

Complexe des accélérateurs du CERN



► H^- (hydrogen anions)

► p (protons)

► ions

► RIBs (Radioactive Ion Beams)

► n (neutrons)


► \bar{p} (antiprotons)

► e^- (electrons)




The detectors at the LHC all measure heavy-ion collisions

A. Andronic

8




LHCb experiment



A detector at the LHC - ALICE

A. Andronic

0




arXiv:1402.4476

ALICE Collaboration: 40 countries, 170 institutions, 2009 members

What are the "control parameters"

- Energy of the collision (per nucleon pair, $\sqrt{s_{NN}}$)
 - Centrality of the collision (number of “participating” nucleons, N_{part})
[at high energies geometric concepts valid: “participant-spectator” picture]
measured in percentage of the geometric cross section ($\sigma_{AB} = \pi(R_A + R_B)^2$)
- NB: we sort the collisions offline, based on detector signals*



...while often taking as reference the measurement in proton-proton collisions (at the same energy), for “hard probes” (pQCD) scaled by the number of collisions corresponding to the given centrality class


What we measure

A. Andronic

11

Production yields and correlations as a function of kinematic quantities:

- transverse momentum, $p_T = p \sin \theta$ (in GeV/c ; recall: $\beta = \frac{v}{c} = \frac{p_c}{E}$)
- rapidity, $y = \frac{1}{2} \ln \frac{E+p_z}{E-p_z} = \tanh^{-1}(p_z/E)$; $p_z = p_L = p \cos \theta$
additive for Lorentz transformations (equivalent of velocity for Galilei)



ALICE, [arXiv:1612.08966](https://arxiv.org/abs/1612.08966)

...poor (wo)man's y : “pseudorapidity”, $\eta = \tanh^{-1}(p_z/p) = \tanh^{-1}(\cos \theta)$
(without particle identification) ... $\eta = y$ for $m = 0$, $\eta \simeq y$ for $p \gg m$

What we measure

We usually measure symmetric collisions of heavy nuclei (Au–Au, Pb–Pb)
pp and p–Pb collisions as reference, but interesting on their own (limits of QGP)
Need many collisions (millions), to sample properly the distributions

- Commonly charged particles, but neutral ones too (via their decays); γ
- The amount of particles (count tracks, assembled from detector points)
- Momentum (via curvature in magnetic field) or energy (in a calorimeter)
...or velocity (via a time-of-flight measurement, resolution 70 ps)
- Identify particles (via energy deposit in detector; ToF; invariant mass)
- Correlations between particles (in each event/collision)


Detector acceptance restricted around midrapidity ($y, \eta = 0$)

Single-particle detection efficiency: about 70-80%
(from detailed Monte Carlo modelling of the detector)

A pictorial summary of collisions at the LHC

A. Andronic

13




©C.Oppedisano/ALICE

Nucleus-nucleus collisions at the LHC

A. Andronic

14




a picture of a central collision (about 3200 primary tracks in $|\eta| < 0.9$); “Camera”: Time Projection Chamber 5 m length, 5 m diam.; 500 mil. pixels; we take 50000 Pb-Pb “pictures” per second (fewer until 2021)

Particle identification

A. Andronic

15




dE/dx : truncated mean of 159 samples along a track; resolution: 5.8%

lines: Bethe-Bloch parametrizations particles and antiparticles are shown

Hadron yields (central collisions)

A. Andronic

16



matter and antimatter produced in equal amounts at the LHC

hadron abundances governed by chemical equilibrium

best fit:

$$T_{CF} = 156.6 \pm 1.7 \text{ MeV}$$

$$\mu_B = 0.7 \pm 3.8 \text{ MeV}$$

$$V_{\Delta y=1} = 4175 \pm 380 \text{ fm}^3$$

$$(\chi^2/N_{df} = 16.7/19)$$

laboratory creation of a piece of hot Universe when $10 \mu\text{s}$ old,
 $T \simeq 10^{12} \text{ K}$

Statistical hadronization (“thermal”) fits

A. Andronic

17

$$n_i = N_i/V = -\frac{T}{V} \frac{\partial \ln Z_i}{\partial \mu} = \frac{g_i}{2\pi^2} \int_0^\infty \frac{p^2 dp}{\exp[(E_i - \mu_i)/T] \pm 1}$$


quantum nr. conservation ensured via chemical potentials:

$$\mu_i = \mu_B B_i + \mu_{I_3} I_{3i} + \mu_S S_i + \mu_C C_i$$

Latest PDG hadron mass spectrum (up to 3 GeV, 600 species)

$$\text{Minimize: } \chi^2 = \sum_i \frac{(N_i^{exp} - N_i^{therm})^2}{\sigma_i^2}$$

N_i : hadron yield $\Rightarrow (T, \mu_B, V)$




Strong (and EM, for Λ) decays play a major role; approx.: $\ln N \sim \ln(mT) - \frac{m}{T}$

Even loosely-bound objects (d , $^3_\Lambda H$) seem produced at $T=156$ MeV (?)

The phase diagram of QCD

A. Andronic

18



Bazavov et al., PLB 795 (2019) 15

Borsanyi et al., PRL 125 (2020) 052001

Andronic et al., Nature 561 (2018) 321

μ_B : a measure of the net-baryon density, or matter-antimatter asymmetry

at LHC, remarkable “coincidence” with Lattice QCD phase crossover

at LHC ($\mu_B \simeq 0$): purely-produced (anti)matter ($m = E/c^2$), as in the Early Universe


$\mu_B > 0$: more matter, from “remnants” of the colliding nuclei

$\mu_B \gtrsim 400$ MeV: *the critical point awaiting discovery (at FAIR?)*

The proton at low resolution ...in collisions at the LHC

A. Andronic


19



The proton at high resolution ...in collisions at the LHC

A. Andronic

20



Strangeness production - from small to large systems

A. Andronic

21

ALICE, [Nature Physics 13 \(2017\) 535](#)

(big geometric) fireball in Pb–Pb reached with violent pp and p–Pb collisions

canonical to grand-canonical strangeness production regime Vislavicius, Kalweit, [arXiv:1610.03001](#)


is the same mechanism at work in small systems (at large multiplicities)?

string hadronization (PYTHIA8) does not describe data

DIPSY (Mueller's dipole, BFKL evolution) fares better Dipole evolution in Impact Parameter Space and rapidity Flensburg, Gustafson, Lonnblad, [arXiv:1103.4321](#)

EPOS LHC: core(QGP)-corona model


new ideas: ropes; thermodynamical string fragmentation



Elliptic flow

A. Andronic

22




$$\frac{dN}{d\varphi} \sim [1 + 2v_1 \cdot \cos(\varphi) + 2v_2 \cdot \cos(2\varphi) + \dots]$$

φ = azimuthal angle with respect to reaction plane

$v_2 = \langle \cos(2\varphi) \rangle$ we call *elliptic flow*


R. Snellings, arXiv:1102.3010



Flow harmonics: all collision systems

A. Andronic

23



Symbols: data, ALICE, PRL 123 (2019) 142301

Lines: hydrodynamics, Schenke, Shen, Tribedy PRC 102 (2020) 044905

Thermodynamic quantities: data-hydrodynamics


A. Andronic

24

Data: effective values
(average over time)
processed through
hydrodynamics

Effective number of
degrees of freedom:
 $g \simeq 30 (\gg 3, \text{ pions})$


ideal gas of quarks
and gluons: $g=52$



Good fluids (ratio viscosity / entropy density)

A. Andronic

25



Bernhard, et al.,

Nature Physics 15 (2019) 1113

Dilute gas: $\eta = \frac{1}{3}n\langle p \rangle \lambda$; $\lambda \sim \frac{1}{n} \rightarrow \eta$ indep. on n

Lowest value of $1/4\pi$ predicted by gravity-gauge duality (AdS-CFT) conjecture


"Simple-minded": $\eta/s \simeq \eta/(k_B n) = p\lambda/3k_B \gtrsim \hbar/3k_B$ ($\Delta p \cdot \Delta x \geq \hbar$)

Flow harmonics


A. Andronic

26

v_2 (elliptic flow) is the dominant flow harmonics, except for most central collisions



Analogy?



http://www.esa.int/Our_Activities/Space_Science/Planck

the non-uniformities (10^{-5}) are the seeds of the
 ~ 100 billion of seen galaxies

Dominant multipole $l \simeq 100$ (angular scale: 1°)


The initial femtoscopy (gluonic) state

A. Andronic

27

central Pb–Pb collision, $\sqrt{s_{NN}}=5.02 \text{ TeV}$ ($R_{\text{Pb}}=6.62 \text{ fm}$)

theoretical calculation (Color Glass Condensate)



Gelis et al, [arXiv:1907.10948](https://arxiv.org/abs/1907.10948)

Mean energy density at the center: 131 GeV/fm^3 ($\alpha_s=0.25$)

Probing early stages

A. Andronic

28


...with "hard probes" ($m \gg T$): jets or high- p_T hadrons (or heavy quarks) produced very early in the collision, $t \simeq 1/m$
(jets - sprays of hadrons from high-speed quarks)

- q, \bar{q}, g travel through QGP, lose energy
- hadronize (neutralize color picking up partners from the vacuum)
- hadrons fly towards detectors

...where we observe a deficit at high momenta (p_T): "jet quenching"
(Bjorken, 1982)

quantified by the nuclear modification factor:




$$R_{AA} = \frac{dN_{AA}/dp_T dy}{N_{coll} \cdot dN_{pp}/dp_T dy}$$



Jets in Pb–Pb collisions

A. Andronic

29



ATLAS

Run: 169045
Event: 1914004
Date: 2010-11-12
Time: 04:11:44 CET

first “direct” observation of “disappearance” of a jet ($E_T > 100$ GeV) in AA

tracks with $p_T > 2.6$ GeV; calorimeter cell thresholds: $E_T > 0.7$ GeV ECAL, $E > 1$ GeV HCAL

quantitatively:

(“leading jet”) $E_{T1} > 100$ GeV vs. the highest E_T jet in the opposite hemisphere, $E_{T2} > 25$ GeV


E_T after subtracting the average contribution of the **underlying event** ($\simeq 75$ GeV for central coll.)

Jets


A. Andronic

30

...in vacuum



...in medium



E : E_T or p_T ; θ_{jet} : opening angle (R or ΔR)


Is this (theoretical expectation) a quantitative effect of the medium?

Jet fragmentation function


A. Andronic

31

Fraction of jet (parton) p_T carried by constituent hadrons



$$z = \frac{p_{T,\text{track}} \cos(r)}{p_{T,\text{jet}}}$$




Significantly modified in Pb-Pb (more high- and low- p_T constituents)

Parton energy loss in the medium (T)

A. Andronic

32


Collisional



Important at low momenta

$$\Delta E \sim L, T^3$$

Radiative



Important at high momenta


$$\Delta E \sim L^2, T^3$$

For energetic partons radiative energy loss dominate, $\Delta E_{rad} \sim \alpha_s \hat{q}$

Jet quenching at the LHC - leading hadrons

A. Andronic

33



measured with "leading hadrons"
 strong suppression, reaching a factor of about 7, $p_T \simeq 7 \text{ GeV}/c$
 remains substantial even at 50-100 GeV/c

not seen with EW observables (γ , W^\pm , Z^0) ... ALICE γ / pQCD NLO calc.
 not seen in p-Pb collisions
 $(p_T \lesssim 3 \text{ GeV}/c$, gluon saturation)

ALICE, JHEP 1811 (2018) 013



a thermal component, $p_T \lesssim 6 \text{ GeV}/c$ (scaling with N_{part}) determined by gluon saturation and collective flow

Jet quenching: transport coefficient

A. Andronic

34

...extracted from a global fit of data



Cao, Wang, arXiv:2002.04028

Apollinario, Lee, Winn, arXiv:2203.16352

Knowledge of the initial state plays a major role

Quark interlude

up to now we only considered hadrons built with *up*, *down*, *strange* quarks
...these are light, masses from a few MeV (*u*, *d*) to \sim 90 MeV (*s*)

what about heavier ones?

...for instance *charm*, which weights about 1.2 GeV

produced in pairs ($c\bar{c}$) in initial hard collisions, $t \sim 1/(2m_c) \leq 0.1 \text{ fm}/c$


preserve their identity throughout the evolution of the fireball

...ideal messengers of the early stage

Charm production in p-Pb, Pb-Pb vs. pp

A. Andronic

36




From charm quarks to hadrons

A. Andronic

37

...an unsolved mystery


- Fragmentation, $c \rightarrow D, D^*, D_s, \Lambda_c$ (empirical functions $D_{c \rightarrow H_c}(z)$)
- Coalescence / (re)combination, $c + q \rightarrow D$, $c + q + q \rightarrow \Lambda_c$
in phase space (in momentum space, resonance recombination)
- Statistical hadronization (as for u,d,s quarks, at T_{CF})



Charm energy loss and flow: data and models


A. Andronic

29



ALI-PUB-501956

ALICE, [arXiv:2110.09420](https://arxiv.org/abs/2110.09420)



In general a good description; a large variation of model results at low p_T
...theoretical uncertainties: nPDFs, hadronization


...some “unification” is being attempted ([arXiv:1803.03824](https://arxiv.org/abs/1803.03824))

Charm spatial diffusion coefficient

A. Andronic

39

key transport parameter (quantifies drag, thermal, recoil forces)



Altenkort et al., PRL 132 (2024) 051902

spatial diffusion coefficient $D_s \simeq T^2/\hat{q}$; $\hat{q} \sim T^3$

Charmonium and deconfined matter

A. Andronic

40

the original idea: Matsui & Satz, [Phys.Lett. B 178 \(1986\) 178](#)

"If high energy heavy-ion collisions lead to the formation of a hot quark-gluon-plasma, then color screening prevents $c\bar{c}$ binding in the deconfined interior of the interaction region."

Refinements: "sequential suppression":

Digal, Satz, Petrecky, [PRD 64 \(2001\) 75](#)


Karsch, Kharzeev, Satz, [PLB 637 \(2006\) 75](#)

no $q\bar{q}$ bound state if

$$r_{q\bar{q}}(T) > r_0(T) \simeq 1/(g(T)T)$$

r_0 Debye length in QGP

$\Rightarrow q\bar{q}$ "thermometer" of QGP




Thermal picture ($n_{partons} = 5.2T^3$ for 3 flavors)

for $T=500$ MeV: $n_p \simeq 84/\text{fm}^3$, mean separation $\bar{r}=0.2$ fm $< r_{J/\psi}$

Charmonium data at RHIC and the LHC

A. Andronic

41



- "suppression" at RHIC (PHENIX)
- dramatically different at the LHC


$$dN_{ch}/d\eta \sim \varepsilon \quad (>20 \text{ GeV/fm}^3, \text{ for } dN_{ch}/d\eta \simeq 2000)$$

...

Charmonium data at RHIC and the LHC

A. Andronic

42



- "suppression" at RHIC (PHENIX)
- dramatically different at the LHC

Statistical Hadronization Model

$$N_{J/\psi} \sim (N_{c\bar{c}}^{dir})^2$$


J/ψ is another observable (charm)
for the phase boundary
calculations are for $T=156$ MeV

$$dN_{ch}/d\eta \sim \varepsilon (>20 \text{ GeV/fm}^3, \text{ for } dN_{ch}/d\eta \simeq 2000)$$

Quark matter in neutron stars?

A. Andronic




43



Core scenarios

A number of possibilities have been suggested for the inner core, including these three options.

- Up quark (u)
- Down quark (d)
- Strange quark (s)
- Anti-down quark (\bar{d})



Quarks

The constituents of protons and neutrons — up and down quarks — roam freely.

Bose-Einstein condensate

Particles such as pions containing an up quark and an anti-down quark combine to form a single quantum-mechanical entity.

Hyperons

Particles called hyperons form. Like protons and neutrons, they contain three quarks but include 'strange' quarks.


©nature

A.Mann, Nature 579 (2020) 20

The quark core in a neutron star ($2M_{Sun}$, $R=12$ km)

A. Andronic

44



Theoretical calculations, Annala et al, [Nature Phys. 16 \(2020\) 907](#)

the closer to ideal gas ($c_s^2 = 1/3$) quark matter is, the larger/heavier the core

NB: not the quark-gluon matter of LHC (antiquarks and gluons largely absent here; neutron star quark matter may be produced at FAIR/GSI)

Summary

- nucleus-nucleus collisions are highly-dynamic
...we establish observables for various stages
- abundance of hadrons with light quarks consistent with chemical equilibrium
the thermal model provides a simple way to access the QCD phase boundary
- charmed hadrons and charmonium also (mostly) produced at phase boundary
... from $c\bar{c}$ pairs (largely) thermalized in QGP
bottomonium suppression observed; bottom thermalization still open
- strong collective flow, well described by hydrodynamics
the derived energy, entropy densities (initial and averaged) are well above the values expected for deconfinement
early (rapid) thermalization remains not fully understood
- we see strong jet quenching (parton energy loss) in quark-gluon matter
data + theoretical models → extraction of transport coefficients

What lies ahead (the next 10 years)

Precision characterization of quark-gluon matter at LHC in Run 3, 4

LHC detectors with major upgrades (ATLAS, CMS major upgrades in LS3)

Interestingly: O-O (and p-O) collisions in 2025 (limits of jet quenching)

Measurements at BNL RHIC (STAR, sPHENIX) and FAIR (CBM)

Extra slides


A. Andronic

QCD EoS

A. Andronic

1


Kojo, Universe 4 (2018) 42, Baym et al., RPP 81 (2018) 056902



Counting charged particles

A. Andronic


2



Nucleus-nucleus collisions: energy density

3

A. Andronic



E_T : transverse energy
(energy built from p_T)

$\varepsilon_{LHC} \simeq 20 - 40 \text{ GeV/fm}^3$
(much above ε_c)

$\varepsilon_{FAIR} \lesssim 1 \text{ GeV/fm}^3$
(around ε_c)

self-similar (Hubble-like) homogeneous (hydrodynamic) expansion of the fireball in the longitudinal (beam) direction
("Bjorken model", 1983)

$$\text{Energy density: } \varepsilon = \frac{1}{A_T} \frac{dE_T}{dy} \frac{1}{c\tau_0}$$


- $A_T = \pi R^2$: transverse area (Pb-Pb: $A_T = 154 \text{ fm}^2$)

- $\tau_0 \simeq 1 \text{ fm}/c$: formation time (establishing the equilibrium) ...not measurable!

Hadron yields at midrapidity (central collisions)

A. Andronic

4



- lots of particles, mostly newly created ($m = E/c^2$)
- a great variety of species:
 - $\pi^\pm (u\bar{d}, d\bar{u})$, $m=140$ MeV
 - $K^\pm (u\bar{s}, \bar{u}s)$, $m=494$ MeV
 - $p (uud)$, $m=938$ MeV
 - $\Lambda (uds)$, $m=1116$ MeV
 - also: $\Xi (dss)$, $\Omega (sss) \dots$
- mass hierarchy in production at high energies ($e^{-m/T}$)

AA, [arXiv:1407.5003](https://arxiv.org/abs/1407.5003)

at lower energies: u, d quarks as "remnants" (stopped) from the incoming nuclei

Viscosity and fluidity

A. Andronic

5




Illustration: Carin Cain

Figure 1: Viscosity—the ratio between a shear force F and the transverse gradient in a velocity profile $\nabla_y v_x$ —causes dissipation in a fluid, which converts part of the kinetic energy in the flow to heat. A "good" fluid is therefore characterized by a small shear viscosity.

$$\frac{F}{A} = \eta \nabla_y v_x, \eta \text{ shear viscosity (in Pa}\cdot\text{s)}$$


Good fluidity: large Reynolds numbers $Re = \frac{mn}{\eta} v L$ (η/n in \hbar)

For relativistic fluids (n not constant): $\eta/(\hbar n) \rightarrow \eta/s$ (in \hbar/k_B)
 s entropy density

Data and hydrodynamics

A. Andronic

6



mass dependence due to collective flow


hydrodynamic models reproduce the data with a very small ratio η/s
viscosity/entropy density, $\eta/s \sim T\lambda c_s$

lower bound conjectured (AdS/CFT): $\eta/s \geq 1/4\pi$ Kovtun, Son, Starinets, [hep-th/0405231](https://arxiv.org/abs/hep-th/0405231)

Elliptic flow: energy dependence

A. Andronic

7



3 regimes:

$v_2 > 0$ at low energies: in-plane, rotation-like emission

$v_2 < 0$ onset of expansion, in competition with shadowing by spectators (which act as a clock for the collective expansion, $t_{\text{pass}} = 40-10$ fm/c)

$v_2 < 0$ at high energies: “free” fireball (almond-shape) expansion (“genuine” elliptic flow)


hydrodynamic description, low η/s

Dileptons: the temperature

A. Andronic

8


...averaged over the space-time history (evolution) of the fireball



The light and the heavy quarks

A. Andronic


9



Nuclear modification: a sketch

A. Andronic


10



Heavy quark energy loss: data and models


A. Andronic

11



b quarks (*non-prompt* D^0) are less suppressed than c quarks (*prompt* D^0)

dead cone effect:



©F.Grosa

$$\Delta E_b < \Delta E_c < \Delta E_{u,d,s}$$

ALICE-PUB-501659


ALICE, [arXiv:2202.00815](https://arxiv.org/abs/2202.00815)

Models describe data ...but not in a perfect agreement

Transport models - schematics

A. Andronic

12



Rapp, Du, [arXiv:1704.07923](https://arxiv.org/abs/1704.07923)

Low p_T : Brownian motion (diffusion) of charm (and b) quarks in QGP

example is for J/ψ , same picture for D , Λ_c (except light quarks 'join' late)

Transport models - in (some) depth

$$\frac{\partial f_Q}{\partial t} = \frac{\partial}{\partial p} \left(A(p)p f_Q + B(p) \frac{\partial f_Q}{\partial p} \right) \simeq \frac{\partial(A(p)p f_Q)}{\partial p} + B \Delta_{\vec{p}} f_Q$$

Fokker-Planck equation for HQ phase-space distribution $f_Q(t, \vec{p})$

special case of Boltzmann eq., for $p^2 \sim m_Q T \gg T^2$ ($m_Q/T \geq 5$)

- A : drag coeff. / thermalization rate; thermal relaxation time $\tau_Q = 1/A(p=0)$
- B momentum diffusion coefficient; *Einstein relation*: $B(p) = A(p)m_Q T$
- Spatial diffusion constant: $D_s = \frac{T}{A(p=0)m_Q}$; $\langle x^2 \rangle - \langle x \rangle^2 = 6D_s t$
- Relation to QCD medium: $D_s(2\pi T) \sim \eta/s(4\pi)$



"Effective" pQCD treatment; no single recipe/formalism, variety of models

$\tau_c \simeq 2 \text{ fm}/c$ (NB: $\simeq 20 \text{ fm}/c$ in leading-order pQCD)

Υ production

A. Andronic

14



CMS, [arXiv:1805.09215](https://arxiv.org/abs/1805.09215)

$\Upsilon(1S)$ suppression to a large extent effect of feed-down from $\Upsilon(2S, 3S)$, which were fully dissociated ("sequential suppression")